



Selectivity of natural, synthetic and environmental estrogens for zebrafish estrogen receptors.

Caroline Pinto, Marina Grimaldi, Abdelhay Boulahtouf, Farzad Pakdel, François Brion, Sélim Aït-Aïssa, Vincent Cavaillès, William Bourguet, Jan-Ake Gustafsson, Maria Bondesson, et al.

► To cite this version:

Caroline Pinto, Marina Grimaldi, Abdelhay Boulahtouf, Farzad Pakdel, François Brion, et al.. Selectivity of natural, synthetic and environmental estrogens for zebrafish estrogen receptors.. *Toxicology and Applied Pharmacology*, 2014, 280 (1), pp.60-69. 10.1016/j.taap.2014.07.020 . inserm-01067804

HAL Id: inserm-01067804

<https://www.hal.inserm.fr/inserm-01067804>

Submitted on 24 Sep 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Selectivity of natural, synthetic and environmental estrogens for zebrafish estrogen receptors

Caroline Pinto^{1*}, Marina Grimaldi^{2*}, Abdelhay Boulahtouf², Farzad Pakdel³, François Brion⁴, Sélim Aït-Aïssa⁴, Vincent Cavaillès², William Bourguet⁵, Jan-Ake Gustafsson^{1,6}, Maria Bondesson¹, Patrick Balaguer^{2#}.

¹Center for Nuclear Receptors and Cell Signaling, Department of Biology and Biochemistry, University of Houston, Houston, Texas 77204-5056, USA. ²Institut de Recherche en Cancérologie de Montpellier, Institut National de la Santé et de la Recherche Médicale U896, Institut Régional de Cancérologie de Montpellier, Université Montpellier 1, 34298 Montpellier, France. ³Institut de Recherche sur la Santé, Environnement et Travail (IRSET), INSERM U1085, Université de Rennes 1, Rennes, France. ⁴Unité Écotoxicologie In Vitro et In Vivo, INERIS, Parc ALATA, 60550 Verneuil-en-Halatte, France. ⁵U1054, Centre de Biochimie Structurale, CNRS UMR5048, Université Montpellier 1 et 2, 34290 Montpellier, France. ⁶Department of Biosciences and Nutrition, Karolinska Institutet, 14183 Huddinge, Sweden

*Equally contributing authors

#To whom correspondence should be addressed.

Patrick Balaguer, Institut de Recherche en Cancérologie de Montpellier, Institut National de la Santé et de la Recherche Médicale U896, Institut Régional de Cancérologie de Montpellier, Université Montpellier 1, Parc Euromédecine, 34298 Montpellier, France.

Email: patrick.balaguer@inserm.fr Tel : 33 4 67 61 24 09 Fax : 33 4 67 61 37 87

Abstract

Zebrafish, *Danio rerio*, is increasingly used as an animal model to study the effects of pharmaceuticals and environmental estrogens. As most of these estrogens have only been tested on human estrogen receptors (ERs), it is necessary to measure their effects on zebrafish ERs. In humans there are two distinct nuclear ERs (hER α and hER β), whereas the zebrafish genome encodes three ERs, zfER α and two zfER β (zfER β 1 and zfER β 2). In this study, we established HeLa-based reporter cell lines stably expressing each of the three zfERs. We first reported that estrogens more efficiently activate the zfERs at 28°C as compared to 37°C, thus reflecting the physiological temperature of zebrafish in wildlife. We then showed significant differences in the ability of agonist and antagonist estrogens to modulate activation of the three zfERs isotypes in comparison to hERs. Environmental compounds (bisphenol A, alkylphenols, mycoestrogens) which are hERs panagonists and hER β selective agonists displayed greater potency for zfER α as compared to zfER β s. Among hER α selective synthetic agonists, PPT did not activate zfER α while 16 α -LE2 was the most zfER α selective compound. Altogether, these results confirm that all hER ligands control in a similar manner the transcriptional activity of zfERs although significant differences in selectivity were observed amongst subtypes. The zfER subtype selective ligands that we identified thus represent new valuable tools to dissect the physiological roles of the different zfERs. Finally, our work also points out that care has to be taken in transposing the results obtained using the zebrafish as a model for human physiopathology.

Keywords: Estrogen receptors, zebrafish, selective estrogens, endocrine disruptors, reporter cell lines

Abbreviations : hERs, human estrogen receptors; zfERs, zebrafish estrogen receptors; AF, transactivation function ; LBD, ligand binding domain; EDCs, endocrine disrupting chemicals; E2, 17 β -Estradiol; 17 α -E2, 17 α -Estradiol ; E1, estrone; E3, estriol; EE2, 17 α -ethynylestradiol; BP2, benzophenone-2; PPT, 4,4',4''-(4-Propyl-[¹H]-pyrazole-1,3,5-triyl) trisphenol; DPN, 2,3-bis(4-hydroxyphenyl)-propionitrile; WAY200070, (7-Bromo-2-(4-hydroxyphenyl)-1,3-benzoxazol-5-ol); FERB033 (2-Chloro-3'-fluoro-3,4'-dihydroxy-[1,1-biphenyl]-4-carboxaldehyde oxime); ERB041, (7-Ethenyl-2-(3-fluoro-4-hydroxyphenyl)-5-benzoxazolol); MPP dihydrochloride, (1,3-Bis(4-hydroxyphenyl)-4-methyl-5--[4-(2-piperidinylethoxy)phenol]-1*H*-pyrazole dihydrochloride); PHTPP, (1,3-Bis(4-hydroxyphenyl)-4-methyl-5--[4-(2-piperidinylethoxy)phenol]-1*H*-pyrazole dihydrochloride); 4OH-Tam, 4-Hydroxytamoxifen; ICI 182,780, (7 α ,17 β -[9[(4,4,5,5,5-pentafluoropentyl)sulfinyl]nonyl]estra-1,3,5(10)-triene-3,17-diol); 16 α -LE2, 3,17-Dihydroxy-19-nor-17 α -pregna-1,3,5(10)-triene-21,16 α -lactone; 8 β -VE2, 8-vinylestra-1,3,5(10)-triene-3,17 β -diol,

Introduction

Estrogens play critical roles in various physiological processes during development and adult life in vertebrates. In mammals, these effects are mediated by members of the nuclear receptor superfamily, the estrogen receptors ER α and ER β (Green et al. 1986; Kuiper et al. 1987). These receptors share a common structural architecture composed of three functional domains. The A/B (or NH₂-terminal domain) is involved in transcriptional activation of gene expression; the C or DNA-binding domain contains a two zinc finger structure, which is important for receptor dimerization and binding of receptors to specific DNA sequences; the E/F or COOH-terminal domain mediates ligand binding, receptor dimerization, nuclear translocation and transactivation of target gene expression in association with coactivators and corepressors (Nilsson et al. 2001). The transactivation functions of both the A/B (AF-1) and the E/F (AF-2) domains are dependent on the cell type and promoter context (Berry et al. 1990). Some ER subtype-selective ligands have been identified, which have different binding affinities for the two estrogen receptors and present variable agonistic or antagonistic characters depending on the ER considered (Delfosse et al. 2012; Escande et al. 2006; Molina-Molina et al. 2008). Dissimilarities in the N-terminal and ligand binding domain (LBD) regions of ER α and ER β explain the differences between the two receptors in their response to various ligands (Matthews et al. 2003; Ogawa et al. 1998).

Even though in mammals only two ER subtypes have been characterized, the presence of three ER subtypes has been reported in teleosts including the zebrafish (*Danio rerio*) (Hawkins et al. 2000; Ma et al. 2000; Menuet et al. 2002). Zebrafish ER α (esr1) is orthologous to the human ER α , while ER β 1 (esr2b) and ER β 2 (esr2a) are orthologs of the human ER β (Bardet et al. 2002). The overall amino-acid sequence identity between the zfER subtypes and their corresponding human ER orthologs is approximately 50% (Menuet et al. 2002). Since these three zfERs are thought to

mediate different biological effects, there is an increased interest in finding subtype-selective zfER ligands.

Xenoestrogens represent a class of endocrine disruptors that affect estrogen signaling. This class of compounds is composed of natural estrogens (phyto and mycoestrogens), and synthetically derived agents including certain pharmaceuticals, pesticides and industrial compounds used in consumer goods (Singleton et al. 2003). Zebrafish is a worldwide recognized vertebrate model to investigate the mode of action of endocrine disrupting chemicals (EDCs) and their reproductive and developmental effects (Segner et al. 2009). Studies on zebrafish have led to significant advances on the effect of EDCs on ER-regulated pathways through the use of zebrafish-specific *in vitro* and *in vivo* models (Menuet et al. 2002; Brion et al. 2012; Cosnefroy et al. 2012; Gorelik et al. 2014). Although all three zfERs subtypes can be activated by 17 β -estradiol (E2), similarly to the mammalian ERs, it has been shown that the capacity of some compounds to transcriptionally activate the estrogen receptors differs between zebrafish and mammals. For instance, the phytoestrogen genistein and the benzophenones BP1, THB and BP2 which are selective human ER β agonists (Escande et al. 2006; Molina-Molina et al. 2008) preferentially activate zfER α (Cosnefroy et al. 2012). However, a detailed analysis of similarities and differences in ligand specificity between zfERs and hERs remains to be performed. The evaluation of the activity and transcriptional profiles of known mammalian estrogenic ligands in the zebrafish model can provide additional information for the analysis of ER-mediated processes in this organism including disruption of these processes by xenoestrogens (Notch and Mayer. 2011).

The purpose of this study was to evaluate the effects of several known natural, environmental and pharmaceutical (anti)estrogenic compounds on the transcriptional activity of the three zfERs and

to compare the data with their activity on hERs. To address this challenge, zfER α , zfER β 1 and zfER β 2 reporter cell lines were established in the same cellular context as we previously used to create hER α and hER β cell lines (Balaguer et al. 1999). In HeLa cells, which stably express an ERE-driven luciferase reporter (HELN cells), we expressed the full-length zfER α , zfER β 1 and zfER β 2, respectively. The resulting HELN-zfERs cell lines are useful tools for the analysis of the effects of estrogenic compounds on gene transactivation by the three zfERs, and also for the comparison of these effects to the results obtained on the hER orthologs. Since zebrafish is used as a model for the study of the effects of xenoestrogens *in vivo*, determining the transcriptional profiles of estrogenic compounds on the zfERs is crucial to support the zebrafish model for ER-related studies and their extrapolation to the mammalian system.

Materials and methods

Materials

Tissue culture plates used in this study came from Greiner Bio-one (Monroe, NC, USA), and media was purchased from Invitrogen (Grand Island, NY, USA). Luciferin (sodium salt) was purchased from Promega (Charbonnières, France). 17 β -Estradiol (E2) (1,3,5 [10]-estratriene-3,17 β -diol), 17 α -Estradiol (17 α -E2) (1,3,5 [10]-estratriene-3,17 α -diol), estrone (E1) (1,3,5[10]-estratriene-3-ol-17-one), estriol (E3) (1,3,5[10]-estratriene-3,16 α ,17 β -triol), 17 α -ethynylestradiol (EE2) (17 α -ethynyl-1,3,5[10]-estratriene-3,17 β -diol), genistein (40,5,7-trihydroxyisoflavone), benzophenone-2 (BP2) (2, 2', 4, 4'-tetrahydroxybenzophenone), α -zearalanol, α -zearalenol, 4-tert-octylphenol and bisphenol A were purchased from Sigma-Aldrich (Saint-Quentin Fallavier, France). 4,4',4''-(4-Propyl-[¹H]-pyrazole-1,3,5-triyl) trisphenol (PPT), 2,3-bis(4-hydroxyphenyl)-propionitrile (DPN), WAY200070 (7-Bromo-2-(4-hydroxyphenyl)-1,3-benzoxazol-5-ol), FERB033 (2-Chloro-3'-fluoro-3,4'-dihydroxy-[1,1-biphenyl]-4-carboxaldehyde oxime), ERB041 (7-Ethenyl-2-(3-fluoro-4-hydroxyphenyl)-5-benzoxazolol), MPP dihydrochloride (1,3-Bis(4-hydroxyphenyl)-4-methyl-5--[4-(2-piperidinyloxy)phenol]-1H-pyrazole dihydrochloride), PHTPP (1,3-Bis(4-hydroxyphenyl)-4-methyl-5-[4-(2-piperidinyloxy)phenol]-1H-pyrazole dihydrochloride) and liquiritigenin were purchased from Tocris Bioscience (Minneapolis, MN, USA). Ferutinin was purchased from Santa Cruz Biotechnology Inc (Dallas, Texas, USA). 4-Hydroxytamoxifen (4OH-Tam) (1-[p-dimethylaminoethoxyphenyl]-1-(4-hydroxyphenyl)-2-phenyl-1-butene), ICI 182,780 (7 α ,17 β -[9[(4,4,5,5,5-pentafluoropentyl)sulfinyl]nonyl]estra-1,3,5(10)-triene-3,17-diol) were obtained from Zeneca (Macclesfield, UK). Raloxifene (6-hydroxy-3-[4-[2-(1piperidinyloxy)phenoxy]-2-(4-hydroxy phenyl)-benzothiophene) came from Eli Lilly (Indianapolis, IN, USA). 3,17-

Dihydroxy-19-nor-17 α -pregna-1,3,5(10)-triene-21,16 α -lactone, named 16 α -LE2 and 8-vinyestra-1,3,5(10)-triene-3,17 β -diol, named 8 β -VE2 is a kind gift of Peter Muhn from Research Laboratories of Schering AG (Berlin, Germany). Compounds were dissolved in dimethyl sulfoxide (DMSO) as 10⁻² M stock solutions and successive dilutions were performed in cell culture medium. The final DMSO concentration never exceeded 0.1% (v/v) of the culture medium.

Cell lines

HELN-zfER α , -zfER β 1 and -zfER β 2 reporter cell lines were established in a similar way as HELN-hER α and -hER β cell lines (Balaguer et al. 1999). Briefly, HELN-zfER α , β 1 and β 2 cell lines cells were obtained by transfection of HELN cells (HeLa cells stably transfected with the ERE- β Globin-Luc-SVNeo plasmid) (Balaguer et al. 1999) by the corresponding pSG5-puro plasmids (pSG5-zfER α -puro, pSG5-zfER β 1-puro and pSG5-zfER β 2-puro, respectively). Cells were selected by geneticin (G418, Sigma-Aldrich) and puromycin (Sigma-Aldrich) at the final concentration of 1 mg/mL and 0.5 μ g/mL, respectively. The best responsive clones were selected based on both EC₅₀ values and luciferase induction factors of E2.

Cell culture conditions

The HELN zfERs and hERs cell lines were cultured in phenol red-free Dulbecco's Modified Eagle's Medium/F12 (DMEM/F12) and supplemented with 5% dextran-coated charcoal-treated fetal bovine serum (FBS-DCC) and 1% antibiotic (penicillin/streptomycin) in a 5% CO₂ humidified atmosphere at 37°C. Selection agents (geneticin 1 mg/ml and puromycin 0.25 μ g/ml)

were maintained in the cell culture medium. Luciferase assays were performed in 5% FBS-DCC medium.

zfER α , zfER β 1 and zfER β 2 transactivation assays

The HELN-zfERs cell lines were seeded at a density of 80,000 cells per well in 96-well white opaque tissue culture plates (Greiner CellStar, USA). Compounds to be tested were added 8 h later, and cells were incubated with compounds at 28°C for 16 h. Cells were maintained in the presence of ligands at 28°C because of the improved transactivation ability of the zfERs at this temperature. Experiments were performed in quadruplicates in at least two independent experiments. At the end of the incubation period, culture medium was replaced with medium containing 0.3 mM luciferin. Luciferase activity was measured for 2 s in intact living cells using a plate reader (PerkinElmer Luminometer, MA, USA). EC50 values were measured using GraphPad Prism software (version 5.04; Graphpad Software Inc., San Diego, CA, USA). hER α and hER β 1 transactivation assays were performed in a similar manner except for the incubation temperature with the ligands (37°C instead of 28°C).

zfERs binding assays

The HELN-zfERs cell lines were seeded at a density of 80,000 cells per well in 96-well white bottom clear tissue culture plates (Greiner CellStar, USA). [³H]-E2 3 nM in the presence or in the absence of non-radioactive E2 10 μ M were added 8 h later, and cells were incubated at 28°C or 37°C for 16h. At the end of the incubation period, unbound material was aspirated and cells

washed three times with PBS. Then, 0.2 ml of LSC-cocktail (Emulsifier-Safe, Packard Bioscience) was added and [^3H]-bound radioactivity was counted. Protein concentrations were measured by Bio-Rad protein assay and used to normalize bound radioactivity values expressed in dpm. Experiments were performed in quadruplicate in at least two independent experiments.

Results

Impact of the temperature on transcriptional activation of zfERs.

Since the functionality of rainbow trout ER has been shown to be temperature-sensitive (Cosnefroy et al. 2009 ; Matthews et al. 2002), we first tested the ability of E2 to activate the zfERs at different temperatures, including a temperature relevant to the physiology of this organism. As seen in Fig. 1A, the EC_{50} value of E2 was lower at 28°C (0.077 nM) than at 31°C (0.266 nM), 34°C (0.433 nM) or 37°C (1.921 nM) for zfER α . Similarly, the EE2 EC_{50} value was lower at 28°C (0.061 nM) than at 31°C (0.127 nM), 34°C (0.19 nM) or 37°C (0.947 nM) (Fig. 1B). EC_{50} of zfER β 1 and zfER β 2 cell lines were also sensitive to the temperature (Fig. 1C and 1D). EC_{50} values for E2 and EE2 were 0.039 and 0.031 nM at 28°C and 0.714 and 0.673 nM at 37°C for zfER β 1. EC_{50} values for E2 and EE2 were approximately 0.118 and 0.031 nM at 28°C and 1.554 and 0.197 nM at 37°C for zfER β 2. As we suspected that the increase in the EC_{50} values was due to zfERs degradation, we measured protein level by saturation ligand binding assay with [^3H -E2]. In HELN-zfER cells, zfER concentration was 8-fold lower at 37°C than 28°C (data not shown).

Effects of human endogenous estrogens on zfERs transcriptional activation

The measured EC50 values for E2 indicates that E2 has a slight zfER β 1 selectivity relative to the other zfER subtypes in this cell context. The reported EC50 values for E2 is approximately 0.017 nM for hER α and 0.068 nM for hER β (Escande et al. 2006), suggesting that E2 has approximately a 6-fold higher potency to transactivate luciferase gene expression in hER α relative to its zebrafish counterpart. 17 α -Ethinylestradiol (EE2), which has a 31-fold greater potency to transactivate the luciferase reporter gene in HELN-hER α compared to hER β (Escande et al. 2006), demonstrated equal agonistic activities on all 3 zfERs with EC values of 0.061 nM for zfER α and 0.031 nM for both zfER β isoforms (Table 1).

We next evaluated the effects of the weak estrogenic stereoisomer of 17 α -estradiol (17 α -E2), estrone (E1) and estriol (E3), on zfER-mediated transcriptional activation (Fig. 2). These endogenous ligands have been shown to behave as full agonists in both of the human ERs with a slight ER α selectivity (Escande et al. 2006 and supplementary data Fig. S1A). The transcriptional profiles obtained by E1, E3 and 17 α -E2 indicate that these estrogens have a stronger potency in transactivating zfER α compared to the zfER β 1 and zfER β 2 isoforms (Fig. 2 A, B, C). E1 and E3 displayed similar potencies on zfER α (EC50 values of 1.7 and 0.55 nM, respectively), zfER β 1 (EC50 values of 5.6 and 2.6 nM, respectively) and zfER β 2 (EC50 values of 12.1 and 8.5 nM, respectively). Among the endogenous estrogens tested, 17 α -E2 showed the weakest agonistic activity with EC50s of 5.3, 24.7 and 45.1 nM for zfER α , zfER β 1 and zfER β 2, respectively.

Selectivity of natural estrogens on zfERs transcriptional activity

We next analyzed the ability of myco and phytoestrogens to activate the zfERs (Fig. 3). The mycoestrogen α -zearalanol (α -zea) exhibited EC₅₀ values of 0.176 nM for zfER α , 0.71 nM for zfER β 1 and 1.2 nM for zfER β 2, which conferred 4-fold and 6.8-fold higher potencies in transactivation assays for HELN-zfER α compared to the zfER β cell lines (Fig. 3A). The other zearalanone analogues (zearalanone, β -zearalanol, zearalenone, α -zearalenol and β -zearalenol) presented a similar profile (zfER α potency > zfER β 1 potency \geq β 2 potency, supplementary data Fig. S2A and data not shown). The phytoestrogen genistein, which present a higher affinity for hER β than for hER α (Escande et al. 2006; Molina-Molina et al. 2008), demonstrated greater affinity for zfER α and zfER β 2 relative to zfER β 1 (Fig. 3B). Genistein activated zfER α and β 2 with a 3.9-4.4 greater potency than zfER β 1 with EC₅₀ values of 40, 50 and 260 nM for zfER α , zfER β 2 and zfER β 1, respectively. Liquiritigenin, another phytoestrogen with higher affinity for hER β than for hER α (supplementary data Fig. S1D), activated zfER α with a 12 greater potency than zfER β 2 with EC₅₀ values of 563 and 6805 nM for zfER α and zfER β 2, respectively, and did not activate zfER β 1 at all (table 1 and Fig. 3C). Finally, ferutin, a phytoestrogen with similar affinity for hER α and hER β but with different maximal activity (Table 1 and supplementary data Fig. S1E) displayed similar maximal activity (100%) but different potency for the zfERs (EC₅₀ values of 8, 3.4 and 132 nM for zfER α , zfER β 1 and zfER β 2, respectively) (Table 1 and Fig 3.D).

Selectivity of environmental compounds on zfERs transcriptional activity

We next analyzed the ability of known environmental estrogenic compounds to activate the zfERs (Fig. 4). Similarly to the human ERs (Delfosse et al. 2012), bisphenol A (BPA) acted as a partial agonist towards the 3 zfERs (Fig. 4A). BPA displayed greater affinity for zfER α than the zfER β subtypes. The EC50 values of BPA were 599 nM, 1.894 μ M and 3.823 μ M for zfER α , zfER β 1 and zfER β 2, respectively (Table 1). The alkylphenol 4-tert-octylphenol (4t-OP) and the nonylphenol mixture (NPm) acted as partial agonists towards zfER α and zfER β 2 and full agonists towards zfER β 1 (supplementary data Fig. 4B and 4C). 4tOP and NPm displayed greater affinity towards zfER α and zfER β 2 relative to zfER β 1. The EC50 values of 4tOP and NPm were 763 and 487 nM for zfER α , 2478 and 4225 nM for zfER β 1 and 1223 and 603 nM for zfER β 2. Benzophenone 2 (BP2), which present a higher affinity for hER β than for hER α (Molina-Molina et al. 2008), demonstrated greater affinity for zfER α and zfER β 2 relative to zfER β 1 (Fig. 4D). BP2 presented a 3-fold higher potency towards zfER α and zfER β 2 compared to zfER β 1 (EC50 values of approximately 251 nM and 233 for zfER α and β 2 respectively, and 803 nM for β 1).

Selectivity of pharmaceuticals on zfERs transcriptional activity

We next evaluated the ability of synthetic ligands to activate transcription through the zfERs. The human ER α selective compound 3,17-dihydroxy-19-nor-17 α -pregna-1,3,5(10)-triene-21,16 α -lactone (16 α -LE2), which has over 1000-fold selectivity in reporter gene activation through hER α than hER β (Escande et al. 2006) also showed greater selectivity for its zebrafish ER α counterpart relative to the zfER β isoforms (Fig. 4A). Among all the compounds tested, 16 α -LE2 displayed the greatest zfER α selectivity with approximately 429 and 115-fold higher potencies to activate

transcription by zfER α compared to zfER β 1 and zfER β 2, respectively. The selective ER α selective compound 4,4',4''-(4-Propyl-[1*H*]-pyrazole-1,3,5--triyl)trisphenol (PPT) did not activate zfER α at all while it partially activated zfER β 1 and zfER β 2 (Fig. 4B). Even though 8-vinylestra-1,3,5(10)-triene-3,17 β -diol (8 β VE2) and 2,3-bis(4-hydroxyphenyl)-propionitrile (DPN) are human ER β selective agonists, these compounds showed better selectivity for zfER α than for the zfER β isoforms (Fig. 4C and 4D). The EC50 values for 8 β VE2 were 32, 188 and 113 nM for zfER α , zfER β 1 and zfER β 2, respectively (Table 1). Likewise, the EC50 values of DPN were 21.7, 238 and 115 nM for zfER α , zfER β 1 and zfER β 2, respectively (Table 1). Similarly, other human ER β selective agonists such as 7-bromo-2-(4-hydroxyphenyl)-1,3-benzoxazol-5-ol (WAY300070), 2-chloro-3'-fluoro-3,4'-dihydroxy-[1,1'-biphenyl]-4-carboxaldehyde oxime (FERB33) and 7-ethenyl-2-(3-fluoro-4-hydroxyphenyl)-5-benzoxazolol (ERB041) displayed greater potencies for zfER α compared to zfER β 1 and zfER β 2 (Table 1 and supplementary data Fig. S2B-D).

Effect of pharmaceutical antagonists on transcriptional activity

The pharmaceutical anti-estrogenic compounds ICI 182,780, raloxifene, 4OH-tamoxifen (4OH-Tam) were tested in the HELN-zfERs and all displayed full antagonistic activity in the three zfER cell lines (Fig. 5A, B, C). No agonistic activity for these three anti-estrogens was observed in our experimental system (data not shown). ICI 182,780 showed slight preferential antagonistic activity towards zfER α (IC50 value of 3 nM) than the zfER β cell lines (IC50 values of 14.7 nM and 28 nM for zfER β 1 and β 2, respectively) (Fig. 5A). 4OH-Tam antagonistic potency was greater for zfER α and zfER β 1 (IC50 values of 8.2 and 4.5nM for zfER α and zfER β 1,

respectively) than for zfER β 2 (IC₅₀ value of 29.4 nM) (Fig. 5B). Raloxifene which has slight preferential antagonistic activity towards hER α (IC₅₀ value of 0.62 nM) than the hER β (IC₅₀ value of 12 nM) was a more powerful antagonist for zfER β 1, with a 4.1 and 9.7-fold greater potencies towards the zfER β 1 subtype compared to zfER α or zfER β 2, respectively (IC₅₀ values of 57, 13.8 and 134 nM for zfER α , zfER β 1 and zfER β 2, respectively) (Fig. 5C).

Another compound tested for its effect on transcriptional inhibition on the HELN-zfERs was the hER α selective agonist and hER β antagonist PPT. As shown in figure 5D, PPT displayed full antagonistic activity towards zfER α (IC₅₀ value of 261 nM) and it was able to partially antagonize estrogenic activation in the zfER β cell lines with IC₅₀ values of 854 and 326 nM for zfER β 1 and zfER β 2, respectively. Finally, antagonists with hER α (MPP) and hER β (PHTPP) selectivity were tested on the HELN-zfERs reporter cells. MPP presented slight zfER β 1 selectivity and PHTPP behaved as zfERs pan-antagonist (Table 1 and supplementary data Fig. S2E-F).

Structural analysis of zfER α selectivity

To gain structural insights into the zfER α selectivity of 16 α -LE₂ in human and zebrafish ERs, we used the web-based server EDMon (Endocrine Disruptor Monitoring; <http://atome2.cbs.cnrs.fr/AT2B/SERVER/EDMon.html>) (Delfosse et al. 2012) to model hER α and hER β in complex with this ligand. The structural basis of the hER α and hER β selectivity towards certain ligands has been associated with two amino acid differences in their ligand binding pockets. Indeed, L384 and M421 of hER α are replaced by M336 and I373 in hER β , respectively (Fig. 7 and Manas et al., 2004). Superimposition of the 16 α -LE₂-bound hER α model

on the crystal structure of hER α in complex with E₂ (PDB code 3UUD) showed that the phenol ring of 16 α -LE₂ occupies the same position as that of E₂ and is engaged in a network of hydrogen bonds with E353 from helix 3 (H3) and R394 from H5 (Fig. 8A). On the other side of the ligand-binding pocket (LBP), it appears that the hydrogen bond observed between the 17-hydroxyl group of E₂ and H524 (H11) is conserved in 16 α -LE₂. The difference between the two complexes resides in the lactone ring of 16 α -LE₂ which points towards M421 (H7) that must undergo a large conformational change to accommodate this additional group. In hER β , the linear M421 is replaced by the branched residue Ileu 373 characterized by a much smaller intrinsic flexibility (Fig. 8B). As a consequence, I373 maintains the synthetic ligand in a position where it interacts unfavorably with M336 (H3). Therefore, 16 α -LE₂ adopts different positions in hER α and hER β , the more constrained environment provided by the latter accounting for the weaker affinity of the ligand for this receptor subtype. The affinity values measured with the zebrafish receptors reflect the variations in the space constraints provided by the different combinations of residues in the three receptor subtypes. With H3 and H7 residues identical to those of the human receptor, zfER α interacts with 16 α -LE₂ with the highest affinity. The slight difference in the binding affinity of 16 α -LE₂ for hER α and zfER α relies most likely on the replacement of L349 (H3) by a methionine residue (M317) (Fig. 7) and a possible loss of a favorable interaction provided by the branched but not by the linear residue (Fig. 8C). With a conserved isoleucine in H7 (I406) and a leucine residue in H3 (L369) (Fig. 7), zfER β 1 displays the most constrained LBP reflecting the weakest binding affinity for 16 α -LE₂. This receptor combines two large residues with low (isoleucine) and medium (leucine) flexibilities (Figure 8D). The replacement of I406 in H7 of zfER β 1 by a leucine residue (L391) (Fig. 7) in zfER β 2 provides a slight gain in LBP plasticity (Figure 8D), in agreement with the slightly better affinity of 16 α -LE₂ for the latter.

The inability of hER β -selective phytoestrogens (genistein and liquiritigenin) and pharmaceuticals (8bv-E2, DPN, WAY300070, FERB033 and ERB041) to activate the zfER β isoforms is explained by the mutation of a critical amino acid involved in genistein binding in hER β . In all zfERs, the position homologous to hER β M336 is occupied, as in hER α , by a leucine residue (Fig. 7)(Sassi-Messai et al. 2009). This amino acid change most likely accounts for the lack of obvious selectivity of the phytoestrogens towards the zfER β subtypes.

Discussion

Screening of endogenous, environmental and synthetic ligands in the HELN-zfER cell lines showed that known mammalian ER ligands are also able to induce transcriptional activity of zebrafish ER subtypes. This study allowed us to assess differences in the potency of the estrogenic compounds among the three zfER subtypes, and compare their selectivity towards hERs using a similar human cellular context. The HELN-zfERs cells were incubated at 28°C after addition of chemicals to the cells because it is a more physiologically relevant temperature for zebrafish, which increased the potency of estradiol approximately 10-fold compared to incubation at 37°C (Fig. 1). Temperature sensitivity of fish ERs has already been reported using reporter gene assays (Cosnefroy et al. 2009; Matthews et al. 2002) and the reason seems to be thermo-dependence of estrogen binding (Matthews et al. 2002; Sumida et al. 2003; Tan et al. 1999). Interestingly, this thermo-dependence is not shared by other zebrafish nuclear receptors like the peroxisome proliferator activated receptor γ (Riu et al. 2014).

Results from this study demonstrated that the selectivity of some of the tested compounds on zfERs differs from that observed between hER α and hER β orthologs. zfER α and zfER β s belong

to the ER α and ER β phylogenetic groups, respectively; however, one to three amino-acids differ in the ligand binding domain of the zfERs compared to their human ER counterparts (Table 3), which certainly explains the differences in selectivity and sensitivity of estrogenic ligands between zebrafish and human ERs.

Among the natural estrogens tested in the HELN-zfER cell lines, 17 β -estradiol (E2) was the compound with the highest potency towards the three zfERs, followed by estriol (E3), estrone (E1) and 17 α -E2. Likewise, a study using transiently-transfected HEK293 cells showed that E2 was the most potent endogenous ligand for zfER α with EC50 of 0.177 nM, very similar to the EC50 value of E2 for zfER α in our study (0.077 nM) (Lange et al. 2012). E2 also is the endogenous compound with the highest affinity for the hERs, and it also displays a better affinity for hER α (Escande et al. 2006). The zfERs were less sensitive to E2 than hERs, with approximately 6 and 2-fold lower transactivation potencies for the zfER α and zfER β subtypes relative to their human orthologs, respectively.

Among the myco and phytoestrogens, α -zearalanol (α -zea) demonstrated the best agonistic activity via the zfER α and zfER β forms with the ability to almost fully activate the luciferase reporter at relative low EC50 values. α -zea preferentially activated zfER α in the HELN-zfERs, which is in accordance with our previous studies performed in zebrafish liver-derived cell lines (Cosnefroy et al. 2012). The phytoestrogen genistein, a full agonist in both human ERs with 10-fold higher selectivity for hER β , showed preferential and equal activity towards zfER α and zfER β 2 over zfER β 1. This phytoestrogen also behaved as a full agonist in the zebrafish cell lines, and displayed slightly higher affinity for zfER α and zfER β 2 than for zfER β 1 (Cosnefroy et al. 2012). Similar transactivation properties by genistein were observed in HeLa cells transiently transfected with the zfERs and an estrogen-dependent luciferase reporter (Sassi-Messai et al.

2009). Lignitrigenin, another phytoestrogen with higher hER β than hER α affinity, also displayed preferential and equal activity towards zfER α and zfER β 2 over zfER β 1. Finally, ferutinin, a terpenoid with similar affinity for hER α and hER β but different activity (full agonist on hER α and partial agonist on hER β) (Ikeda et al. 2002), behaved as a full agonist on the three zfERs and displayed slightly higher affinity for zfER α and zfER β 1 than for zfER β 2.

Among environmental pollutants, bisphenol A (BPA) also presented higher affinity for zfER α relative to the zfER β subtypes. Accordingly, in the zebrafish-derived cell lines (Cosnefroy et al. 2012), BPA also displayed better potency towards zfER α . Considering reporter gene activation in the human ERs, BPA displays similar agonistic properties between hER α and hER β , presenting slightly better efficacy towards hER β (Delfosse et al. 2012). The alkylphenols 4tOP and NPM also showed better potency towards zfER α than the zfER β subtypes. The benzophenone derivative BP2 behaved as a full agonist in the three zfERs, with similar EC50s for both zfER α and zfER β 2. Cosnefroy et al. (2012) also found that BP2 has a slightly higher activity towards zfER α , with approximately 1.36 and 2 fold- higher potencies compared to zfER β 2 and β 1, respectively (Cosnefroy et al. 2012).

17 α -EE2 was the pharmaceutical compound that displayed the highest affinity towards the zfERs, with almost equivalent potencies in transactivating the three zfER subtypes. Similarly to hERs, 16 α -E2, which has 1000-fold selectivity for hER α (Escande et al. 2006), also exhibited higher affinity for zfER α compared to the zfER β subtypes (approximately 100 and 370-fold higher potencies than zfER β 2 and zfER β 1, respectively) and it is the most selective compound for zfER α found in this chemical screening. Conversely, PPT, another hER α selective agonist that is unable to activate hER β (Escande et al. 2006), was devoid of agonistic activity in the HELN-zfER α cells, and instead behaved as an antagonist to zfER α , while it slightly activated the zfER β

subtypes. IC50 values obtained in zfERs cells indicated that PPT displayed preferential affinity for zfER α than for zfER β (Table 2 and Fig. 7). Similarly to phytoestrogens, known human ER β selective synthetic agonists, such as DPN, 8 β -VE2, FERB033, ERB033 and WAY200070 preferentially activated zfER α .

A recent study reported differences in ER subtype tissue localization in the developing zf larvae: zfER α is selectively expressed in the heart valves, zfER β 2 is robustly expressed in the liver, whereas zfER β 1 is not detected (Gorelik et al. 2014). In this same study, treatment of a transgenic zebrafish model containing a 5x consensus ERE sequence upstream of a GFP reporter (5xERE:GFP) with genistein, BPA and DPN preferentially induced GFP expression in the heart valves. PPT, on the other hand, showed GFP labeling of only the larval liver. Our study confirms the *in vivo* findings that these compounds present opposite selectivity of what has been observed for the human ER subtypes. Moreover, as the selectivity of several ligands is not conserved between human and zf ER subtypes, the determination of the transcriptional profile of estrogenic compounds towards the zfERs is crucial before their use *in vivo* to help elucidate distinct roles of each ER subtype in zf.

The hERs pan-antagonists ICI 182,780, 4-OH tamoxifen all acted as pure antagonists in the three zfERs. ICI 182,780 presented slight zfER α selectivity while 4-OH tamoxifen antagonized more efficiently zfER β 1 and zfER α than zfER β 2. In accordance with our results, ICI 182,780 blocks EE2-mediated transcriptional activation with all three isoforms of the zfERs in MDA-MB-231 cells transiently transfected with the zfERs (Notch and Mayer ; 2011). *In vivo*, treatment of zebrafish larvae with either ICI 182,780 or 4-OH tamoxifen reduce the E2-induced fluorescence in liver of transgenic 5XERE:GFP zebrafish (Gorelik and Halpern. 2011). Although ICI 182,780 has a higher antagonistic activity towards zfER α compared to zfER β 2 and β 1, ICI 182,780 is also

efficient in reducing or blocking the GFP induced by various estrogens and pro-estrogens in the brain of cyp19a1b-GFP larvae (Brion et al. 2012). The hER α (raloxifen, MPP) and hER β selective (PHTPP) antagonists were also tested on the HELN-zfERs reporter cells. Slight zfER α selectivity was also observed for PHTPP whereas raloxifen and MPP antagonized more efficiently zfER β 1 than zfER α and zfER β 2. Thus, there is also a non-conserved selectivity between zfERs and their human orthologs for antiestrogens.

To conclude, we have shown here that there are clear differences between the selectivity of various (anti)estrogens for zebrafish and human ER isoforms, establishing the fact that a direct translation of (anti)estrogenic effects (activities or potencies) from mammals to zebrafish is not possible. Although none of the tested compounds specifically activated either zebrafish or human ERs, the differences revealed in this study in terms of transcriptional activities towards human and zebrafish ERs highlight the need to take into account i) the species of origin and ii) the ER subtype when assessing the estrogenic potency of chemicals. This is particularly important in regards to EDCs screening for hazard assessment, since at the present time established test guidelines are only based on human cell lines expressing human estrogen receptors. To this end, such newly established *in vitro* cell lines together with those previously established in ZFL cells (Cosnefroy et al. 2012) can serve as useful screening tools to address estrogenic potency of chemicals for piscine models. Hence, an initial screening should be followed up with an ER-subtype specific analysis using both human and zebrafish ERs to elucidate the full spectrum of ER-mediated xenoestrogens effects.

Acknowledgements

This study was funded by grants from Agence Nationale de la Recherche Contaminants, Ecosystèmes, Santé Project “BISCOT” 2010, by the P189 NEMO project financed by the French Ministry of Environment, the US-Environmental Protection Agency (grant number R834289), the Texas Emerging Technology Fund under Agreement 300-9-1958, the National Institutes of Health’s National Institute of Environmental Health Sciences (grant number R21ES020036).

We acknowledge Peter Muhn (Schering AG) for the kind gift of 16 α -LE2 and 8 β -VE2.

References

- Balaguer P, Francois F, Comunale F, Fenet H, Boussioux AM, Pons M, et al. Reporter cell lines to study the estrogenic effects of xenoestrogens. *The Science of the total environment*. 1999;233:47-56.
- Bardet PL, Horard B, Robinson-Rechavi M, Laudet V, Vanacker JM. Characterization of oestrogen receptors in zebrafish (*Danio rerio*). *Journal of molecular endocrinology*. 2002;28:153-63.
- Berry M, Metzger D, Chambon P. Role of the two activating domains of the oestrogen receptor in the cell-type and promoter-context dependent agonistic activity of the anti-oestrogen 4-hydroxytamoxifen. *The EMBO journal*. 1990;9:2811-8.
- Brion F, Le Page Y, Piccini B, Cardoso O, Tong SK, Chung BC, et al. Screening estrogenic activities of chemicals or mixtures in vivo using transgenic (cyp19a1b-GFP) zebrafish embryos. *PLoS one*. 2012;7:e36069.
- Cosnefroy A, Brion F, Guillet B, Laville N, Porcher JM, Balaguer P, et al. A stable fish reporter cell line to study estrogen receptor transactivation by environmental (xeno)estrogens. *Toxicology in vitro*. 2009;23:1450-4.
- Cosnefroy A, Brion F, Maillot-Marechal E, Porcher JM, Pakdel F, Balaguer P, et al. Selective activation of zebrafish estrogen receptor subtypes by chemicals by using stable reporter gene assay developed in a zebrafish liver cell line. *Toxicological sciences*. 2012;125:439-49.
- Delfosse V, Grimaldi M, Pons JL, Boulahtouf A, le Maire A, Cavailles V, et al. Structural and mechanistic insights into bisphenols action provide guidelines for risk assessment and discovery of bisphenol A substitutes. *Proceedings of the National Academy of Sciences of the United States of America*. 2012;109:14930-5.
- Escande A, Pillon A, Servant N, Cravedi JP, Larrea F, Muhn P, et al. Evaluation of ligand selectivity using reporter cell lines stably expressing estrogen receptor alpha or beta. *Biochemical pharmacology*. 2006;71:1459-69.
- Green S, Kumar V, Krust A, Walter P, Chambon P. Structural and functional domains of the estrogen receptor. *Cold Spring Harbor symposia on quantitative biology*. 1986;51:751-8.
- Gorelick DA, Halpern ME. Visualization of estrogen receptor transcriptional activation in zebrafish. *Endocrinology*. 2011;152:2690-703.
- Gorelick DA, Iwanowicz LR, Hung AL, Blazer VS, Halpern ME. Transgenic Zebrafish Reveal Tissue-Specific Differences in Estrogen Signaling in Response to Environmental Water Samples. *Environmental health perspectives*. 2014;122:356-62.
- Hawkins MB, Thornton JW, Crews D, Skipper JK, Dotte A, Thomas P. Identification of a third distinct estrogen receptor and reclassification of estrogen receptors in teleosts. *Proceedings of the National Academy of Sciences of the United States of America*. 2000;97:10751-6.
- Ikeda K, Arao Y, Otsuka H, Nomoto S, Horiguchi H, Kato S, Kayama F. Terpenoids found in the umbelliferae family act as agonists/antagonists for ERalpha and ERbeta: differential transcription activity between ferutinine-liganded ERalpha and ERbeta. *Biochem Biophys Res Commun*. 2002; 291:354-60.
- Kuiper GG, Enmark E, Pelto-Huikko M, Nilsson S, Gustafsson JA. Cloning of a novel receptor expressed in rat prostate and ovary. *Proceedings of the National Academy of Sciences of the United States of America*. 1996;93:5925-30.
- Lange A, Katsu Y, Miyagawa S, Ogino Y, Urushitani H, Kobayashi T, et al. Comparative responsiveness to natural and synthetic estrogens of fish species commonly used in the laboratory and field monitoring. *Aquat Toxicol*. 2012;109:250-8.

- Ma CH, Dong KW, Yu KL. cDNA cloning and expression of a novel estrogen receptor beta-subtype in goldfish (*Carassius auratus*). *Biochimica et biophysica acta*. 2000;1490:145-52.
- Manas ES, Unwalla RJ, Xu ZB, Malamas MS, Miller CP, Harris HA, Hsiao C, Akopian T, Hum WT, Malakian K, Wolfrom S, Bapat A, Bhat RA, Stahl ML, Somers WS, Alvarez JC. Structure-based design of estrogen receptor-beta selective ligands. *J Am Chem Soc*. 2004;126:15106-19.
- Matthews J, Gustafsson JA. Estrogen signaling: a subtle balance between ER alpha and ER beta. *Molecular interventions*. 2003;3:281-92.
- Matthews JB, Fertuck KC, Celius T, Huang YW, Fong CJ, Zacharewski TR. Ability of structurally diverse natural products and synthetic chemicals to induce gene expression mediated by estrogen receptors from various species. *The Journal of steroid biochemistry and molecular biology*. 2002;82:181-94.
- Menuet A, Pellegrini E, Anglade I, Blaise O, Laudet V, Kah O, et al. Molecular characterization of three estrogen receptor forms in zebrafish: binding characteristics, transactivation properties, and tissue distributions. *Biology of reproduction*. 2002;66:1881-92.
- Molina-Molina JM, Escande A, Pillon A, Gomez E, Pakdel F, Cavailles V, et al. Profiling of benzophenone derivatives using fish and human estrogen receptor-specific in vitro bioassays. *Toxicology and applied pharmacology*. 2008;232:384-95.
- Nilsson S, Makela S, Treuter E, Tujague M, Thomsen J, Andersson G, et al. Mechanisms of estrogen action. *Physiological reviews*. 2001;81:1535-65.
- Notch EG, Mayer GD. Efficacy of pharmacological estrogen receptor antagonists in blocking activation of zebrafish estrogen receptors. *General and comparative endocrinology*. 2011;173:183-9.
- Ogawa S, Inoue S, Watanabe T, Hiroi H, Orimo A, Hosoi T, et al. The complete primary structure of human estrogen receptor beta (hER beta) and its heterodimerization with ER alpha in vivo and in vitro. *Biochemical and biophysical research communications*. 1998;243:122-6.
- Riu A, McCollum CW, Pinto CL, Grimaldi M, Hillenweck A, Perdu E, et al. Halogenated Bisphenol A analogs act as obesogens in zebrafish larvae (*Danio rerio*). *Toxicological sciences*. 2014; In Press.
- Sassi-Messai S, Gibert Y, Bernard L, Nishio S, Ferri Lagneau KF, Molina J, et al. The phytoestrogen genistein affects zebrafish development through two different pathways. *PloS one*. 2009;4:e4935.
- Segner H. Zebrafish (*Danio rerio*) as a model organism for investigating endocrine disruption. *Comparative biochemistry and physiology Toxicology & pharmacology*. 2009;149:187-95.
- Singleton DW, Khan SA. Xenoestrogen exposure and mechanisms of endocrine disruption. *Frontiers in bioscience : a journal and virtual library*. 2003;8:110-8.
- Sumida K, Ooe N, Saito K, Kaneko H. Limited species differences in estrogen receptor alpha-mediated reporter gene transactivation by xenoestrogens. *The Journal of steroid biochemistry and molecular biology*. 2003;84:33-40.
- Tan NS, Frecer V, Lam TJ, Ding JL. Temperature dependence of estrogen binding: importance of a subzone in the ligand binding domain of a novel piscine estrogen receptor. *Biochimica et biophysica acta*. 1999;1452:103-20.

Figure legends

Figure 1. Zebrafish ERs are thermosensitive. Transcriptional activity of zfER α in response to different concentrations of E2 (A) and EE2 (B) was measured at 28, 31, 34 and 37°C. Transcriptional activity of zfER β 1 (C) and zfER β 2 (D) in response to E2 and EE2 were measured at 28 and 37°C. Results are expressed as % 10 nM E2. Standard deviations are from independent experiments done in quadruplicate.

Figure 2. Transcriptional activity of zfER α , zfER β 1 and zfER β 2 in response to endogenous estrogens. HELN-zfER α (■), HELN zf-ER β 1 (▲), HELN-zfER β 2 (○) cells were exposed to different concentrations of E1, E3 and 17 α E2. Results are expressed as % of 10 nM E2 treatment. Standard deviations from at least 2 independent experiments done in quadruplicate are indicated.

Figure 3. Transcriptional activity of zfER α , zfER β 1 and zfER β 2 in response to myco and phytoestrogens. HELN-zfER α (■), HELN zf-ER β 1 (▲), HELN-zfER β 2 (○) cells were exposed to different concentrations of α -zea, genistein, liquiritigenin and ferutinin. Results are expressed as % of 10 nM E2 treatment. Standard deviations from at least 2 independent experiments done in quadruplicate are indicated.

Figure 4. Transcriptional activity of zfER α , zfER β 1 and zfER β 2 in response to environmental estrogens. HELN-zfER α (■), HELN zf-ER β 1 (▲), HELN-zfER β 2 (○) cells were exposed to different concentrations of BPA, 4-tet-octylphenol, NPm and BP2. Results are expressed as % of 10 nM E2 treatment. Standard deviations from at least 2 independent experiments done in quadruplicate are indicated.

Figure 5. Transcriptional activity of zfER α , zfER β 1 and zfER β 2 in response to pharmaceuticals. HELN-zfER α (■), HELN zf-ER β 1 (▲), HELN-zfER β 2 (○) cells were exposed to different concentrations of 16 α LE2, PPT, 8 β VE2 and DPN. Results are expressed as % of 10 nM E2 treatment. Standard deviations from at least 2 independent experiments done in quadruplicate are indicated.

Figure 6. Transcriptional activity of zfER α , zfER β 1 and zfER β 2 in response to synthetic antagonists. HELN-zfER α (■), HELN zf-ER β 1 (▲), HELN-zfER β 2 (○) cells were exposed to different concentrations of ICI 182,780, 4OH-Tamoxifen, raloxifen and PPT together with 1 nM E2. Results are expressed as % of E2 treatment. Standard deviations from at least 2 independent experiments done in quadruplicate are indicated.

Figure 7. Amino acid alignment of human ER β and zebrafish ERs ligand binding domains (LBDs) with par of the human ER α LBD (Met 342 to Ileu 426). The three amino acids involved in the binding of estrogens which differed between ERs are framed. Numbers above the human sequences indicate the number of amino acid residue within the total human ER α and ER β sequences. The GenBank accession numbers of the ER sequences in this alignment are : human ER α NP_000116; human ER β NP_001428; zebrafish ER α NP_694491; zebrafish ER β 1 CAC93848.1 and zebrafish ER β 2 CAC93849.1.

Figure 8. Modeling of the interaction between 16 α -LE₂ and the human (h) and zebrafish (zf) estrogen receptors. Superposition of the structures of hER α (A) and hER β (B) LBDs bound to E₂ (green) onto the molecular models of the 16 α -LE₂-bound hER α (A) and 16 α -LE₂-bound hER β (B) LBDs (yellow). In hER α , the lactone ring of 16 α -LE₂ points towards M421 (H7) which undergoes a large conformational change (black arrow) to accommodate this additional group. In

hER β , the linear M421 is replaced by the branched residue I373 which is characterized by a much smaller intrinsic flexibility that maintains the synthetic ligand in a position (black arrow) where it interacts unfavorably with M336 (H3, red asterisk). Key hydrogen bonds between the protein and the ligands are highlighted by dotted lines. W denote a water molecule. Superposition of the structures of hER α (C) and hER β (D) LBDs bound to E₂ (green) onto the molecular models of the 16 α -LE₂-bound zfER α (C) and 16 α -LE₂-bound zfER β 1 or zfER β 2 (yellow and violet, respectively) (D) LBDs. The steric constraints observed in hERs also apply to zfERs and account for the differential affinity of 16 α -LE₂ for the zfER subtypes. In (C) the dotted line highlights a stabilizing interaction between hER α and the ligand that is probably missing in zfER α .




Figure 1




Figure 2




Figure 3





Figure 4






Figure 6

	349		384		421
hER α	MMGLLTNL	ADRELVHMINWAKRVP	GFVDLT	LHDQVHLL	ECAWLEILMIGLVWRSMEHPGKLLFAPNLLDRNQGKCVEGMVEI
zfER α	MMSLLTNM	ADKELVHMI	AWAKKVP	GFQDLSLHDQVQLLESSWLEVL	MIGLIWRSIHSPGKLIFAQDLILDRSEGECEGMAEI
	301		336		373
hER β	SLTKL	ADKELVHMISWAKKIP	GFVELSLFDQVR	LLESCWMEVLM	MGLMWRSIDHPGKLIFAPDLVDRDEGKCVEGILEI
zfER β 1	SLTSL	ADKELVLMISWAKKIP	GFVELTSLDQVHLL	ECCWLDIILMGLMWRSVDHPGKLIFTPDLKLNREEGNCVEGIMEI	
zfER β 2	SLTNL	ADKELVHMISWAKKIP	GFVELSLFDQVHLL	ECCWLEVL	MGLMWRSVNHHPGKLIFSPDLSLSRDESSCVQGLVEI

Figure 7




Figure 8